


Trends in influenza occurrence in the 2020/21 season in Japan	242	Global influenza activity in 2021	259
Monitoring influenza activity by ILI surveillance, 2020-2021, Mie Prefecture.....	243	An epidemic of RS virus infection, Tokyo Prefecture: description and genetic analysis week 1 to week 31, 2021.....	261
An influenza outbreak during the 2020/21 season in Yamagata Prefecture.....	246	Two COVID-19 clusters where identifying close contacts was difficult at the time of initial detection	263
Analysis of influenza viruses isolated in the 2020/21 season, Japan	247	Analysis of replacement of SARS-CoV-2 strains with a variant carrying the L452R mutation in Japan	265
Influenza vaccination coverage and seroprevalence of influenza, Japan-FY 2020, National Epidemiological Surveillance of Vaccine-preventable Diseases	252	Epidemiological and molecular characteristics of the SARS-CoV-2 Delta variant cases in the early epidemic stage in Japan	267
Effectiveness of influenza vaccine among children younger than 3 years: summary results of the 2018/19 and 2019/20 influenza seasons.....	255	Information on HIV/AIDS cases under the Infectious Diseases Control Law (the National AIDS Surveillance Committee, August 24, 2021).....	269
Update on avian and swine influenza virus infections in humans.....	257	Pathogens detected in confirmed and suspected COVID-19 cases reported in the NESID Infectious Agent Surveillance System (January 2020 to November 2021)	270

<THE TOPIC OF THIS MONTH>
Influenza 2020/21 season, Japan

Figure 1. Weekly number of isolated influenza viruses and reported influenza patients/sentinel sites, week 36 of 2018 to week 35 of 2021, Japan



*Indicates the nationwide start of influenza season: 1.0 patient/week

In the 2020/21 season (week 36 in September 2020 to week 35 in August 2021), the number of reported influenza cases did not increase during the usual epidemic period of the calendar year. Multiple other indicators also remained at remarkably low levels, and no signs suggestive of an influenza epidemic were observed. This may have been due to the epidemic of coronavirus disease 2019 (COVID-19) and countermeasures against it.

Epidemiology of the 2020/21 influenza season: Under the National Epidemiological Surveillance of Infectious Diseases (NESID) system, approximately 5,000 influenza sentinel sites nationwide (approximately 3,000 pediatric and 2,000 internal medicine sentinel sites) report patients diagnosed with influenza on a weekly basis. In the 2020/21 season, the maximum number of reported influenza cases per sentinel was 0.02/week (98 patients reported/week) in the fifth week of 2021, and no week exceeded 1.00/week, which is the threshold for the nationwide start of the influenza season (Fig. 1) (see p. 242 of this issue). Based on the reported sentinel surveillance data, the estimated cumulative total number of medically-attended influenza patients nationally was approximately 14,000 (week 36 of 2020 to week 17 of 2021), which was substantially lower than that during the same period of the past two seasons (7,289,000 patients in the 2019/20 season and 12,005,000 patients in the 2018/19 season). The total number of hospitalized patients in the hospitalized influenza patient surveillance (data from approximately 500 designated sentinel hospitals with at least 300 beds) was also extremely low at 131 patients in the 2020/21 season (week 36 of 2020 to week 17 of 2021), compared with 20,719 patients in the 2018/19 season and 13,011 patients in the 2019/20 season. The number of schools with class closures reported to the "influenza-like illness surveillance" system (school closure and absenteeism surveillance) was 3 schools (week 36 of 2020 to week 9 of 2021), a drastic decrease from the same period of the last two seasons when more than 20,000 schools with class closures were reported. Among patients diagnosed with acute encephalitis (including encephalopathy), a category V notifiable infectious disease, there were no cases diagnosed as influenza encephalopathy. No signs of an influenza epidemic were observed through the influenza-like illness surveillance system conducted in Mie Prefecture (see p.243 of this issue).

Isolation/detection of influenza virus in the 2020/21 season: Prefectural and municipal public health institutes reported the isolation/detection of 7 influenza viruses in the 2020/21 season (Table 1 on p.241). Among them, 5 were reported from influenza sentinel sites and 2 from non-sentinel sites (Table 2 on p.241). Regarding isolates/detections of influenza, two subtype A/H1pdm09 viruses and four subtype A/H3 viruses were reported, but there were no reports of type B virus for either the Victoria or the Yamagata lineage. One type C influenza virus was detected (Table 2 on p.241). Two subtype A/H1pdm09 viruses were reported from Nagasaki Prefecture in weeks 43 and 44 of 2020, two subtype A/H3 viruses were reported from Yamagata Prefecture in week 6, and another two subtype A/H3 viruses were reported from Miyagi Prefecture in week 9 of 2021 (Figs. 1 and 2 on p.241).

Genetic and antigenic characteristics of the 2020/21 isolates: The National Institute of Infectious Diseases (NIID) conducted genetic and antigenic analyses of isolates from Japan and other Asian countries. Antigenic analyses were done using ferret antiserum (see p. 247 of this issue).

The two subtype A/H1pdm09 strains detected in Japan were classified into genetic subclade 183P-5A1 (D187A, Q189E) based on

(THE TOPIC OF THIS MONTH-Continued)

the phylogenetic tree analysis of the hemagglutinin (HA) gene. Antigenic analysis revealed that the two isolates reacted well with ferret antiserum against the “egg-isolated” A/Guangdong-Maonan/SWL1536/2019 (183P-5A1), the vaccine strain for the 2020/21 season.

For subtype A/H3, phylogenetic analysis of the HA gene of four domestic strains and 14 isolates from Laos and Nepal showed that the domestic and Laotian strains belonged to subclade 3C.2a1b.2a1 and the Nepalese strain belonged to subclade 3C.2a1b.2a2. Antigenic analysis showed that all tested strains had poor reactivity with ferret antiserum against the “cell culture propagated” A/Hong Kong/45/2019 and “egg-isolated” A/Hong Kong/2671/2019, the vaccine strains for the 2020/21 season that belong to subclade 3C.2a1b.1b.

Although the type B/Victoria lineage virus was not detected in Japan, the HA gene phylogenetic tree analysis of the strains detected overseas showed that most of the strains belonged to subclade 1A.3a1 or 1A.3a2. In antigenic analysis, these viruses did not react well with ferret antiserum against “cell culture propagated” or “egg isolated” B/Washington/02/2019, the vaccine strain for the 2020/21 season. No viruses of the type B/Yamagata lineage were analyzed.

Resistance to antivirals among the 2020/21 isolates: The two subtype A/H1pdm09 viruses isolated in Japan during the 2020/21 season were not resistant to neuraminidase (NA) inhibitors (oseltamivir, peramivir, zanamivir, and laninamivir) or a cap-dependent endonuclease inhibitor (baloxavir), but were resistant to amantadine. The 14 subtype A/H3 viruses isolated in Japan and overseas (Nepal and Laos) were not resistant to NA inhibitors or a cap-dependent endonuclease inhibitor, but were resistant to amantadine. No strains of type B virus were available for testing (see p.247 of this issue).

Seroprevalence among the Japanese population prior to the 2020/21 season: Based on seroprevalence surveys conducted under the National Epidemiological Surveillance of Vaccine-Preventable Diseases program, seroprevalence levels were measured from 3,244 serum samples collected between July to September 2020 (see p.252 of this issue). The age groups with the highest seroprevalence (measured as HI antibody-positive, titer $\geq 1:40$) were 10-24-year-olds ($\geq 40\%$) for A/Guangdong-Maonan/SWL1536/2019 [subtype A/H1pdm09 vaccine strain], 5-49- and ≥ 70 -year-olds (41-83%) for A/Hong Kong/2671/2019 [subtype A/H3 vaccine strain], 15-39-year-olds ($\geq 60\%$) for B/Phuket/3073/2013 [B/Yamagata lineage vaccine strain], and 40-54-year-olds (30-40%) for B/Victoria/705/2018 [B/Victoria lineage vaccine strain]. Antibody prevalence against A/Hong Kong/2671/2019 [A/H3 subtype vaccine strain] was the highest among the four strains analyzed.

Influenza vaccine: The tetravalent vaccine for the 2020/21 season contained antigens of two type A and two type B strains. Approximately 33,420,000 vial-equivalent doses (estimated on the assumption that 1 vial contained 1 mL) were produced, of which an estimated 32,740,000 vials were used.

For the 2021/22 season, A/Victoria/1/2020 (IVR-217) was selected as the vaccine strain for subtype A/H1pdm09 and A/Tasmania/503/2020 (IVR-221) as the vaccine strain for subtype A/H3. For the vaccine strain of the B/Yamagata lineage, B/Phuket/3073/2013 was selected, and B/Victoria/705/2018 (BVR-11) was selected as the vaccine strain for the B/Victoria lineage (https://www.mhlw.go.jp/stf/shingi2/0000203023_00005.html).

A multicenter case-control study on the effectiveness of two doses of the influenza vaccine in children under three years of age was conducted, and two doses of vaccine were considered to reduce the risk of symptomatic influenza by about half (see p.255 of this issue).

Human infection with avian and swine influenza virus: Since September 2020, several human cases of subtype A/H5 influenza virus infection have been reported. Human cases of highly pathogenic avian influenza (HPAI) A (H5N1) virus infection have been reported from Laos and India, and human cases of A(H5N6) virus infection have been confirmed in China and Laos. In December 2020, the first human case of A(H5N8) virus infection was also confirmed in Russia. With the addition of these cases, the total number of human cases of influenza subtype A/H5 virus infection has reached 863 cases (including 456 fatal cases) since 2003. Human cases of A(H7N9) virus infection have been confirmed in 1,568 cases (including 616 deaths) since 2013, but there have been no reports since March 2019. Human cases of A(H9N2) virus infection have been confirmed in China and Cambodia since October 2020, but all cases were mild. In April 2021, the first human case of A(H10N3) virus infection was reported in China. In addition, HPAI due to subtype A/H5 viruses have occurred among poultry and wild birds in Asia, including Japan, Europe, and Africa, since September 2020.

Swine may be an intermediate “mixing vessel” host, generating reassortments of mammalian and avian influenza viruses. During the 2020/21 season, human cases of A(H3N2)v, A(H1N1)v, and A(H1N2)v virus infections were reported in the United States and Canada, and since October 2020, human cases of swine influenza viruses were also reported from countries such as Australia, China, Europe, and Brazil. Sporadic cases of human infection with avian influenza viruses have been reported in neighboring countries of Japan, and reassortant viruses of human A/H1pdm09 and swine influenza viruses have been detected in swine in Japan. It is necessary to monitor the occurrence of these viruses closely (see p.257 of this issue).

Conclusion: Since the 19th week of 2020, the number of reported influenza cases decreased and remained at a low level even during the regular influenza season in the 2020/21 season. Furthermore, there were no signs suggestive of an influenza epidemic based on multiple indicators. On the other hand, from the summer of 2021 onward, the number of reported influenza cases has been increasing overseas, suggesting the possibility that influenza activity may increase in the 2021/22 season (see p.259 of this issue). In preparation for the resurgence of influenza, it is important to continue public health measures such as vaccination of high-risk groups including the elderly, together with comprehensive monitoring of influenza, such as year-round isolation and detection of influenza viruses, analyses of the antigenic and genetic mutations of circulating strains, and surveys of the emergence of drug-resistant strains and seroprevalence levels in the population.

Note: For IASR reporting, influenza nomenclature is based on the virus information available. Influenza viruses are classified by type, subtype, lineage, and strain based on hemagglutination (HA), neuraminidase (NA), and other information including nucleotide sequence:

- When both HA and NA typing have been performed, the names are listed fully [e.g., A(H1N1)pdm09, A(H3N2), A(H5N1)].
- When NA typing has not been performed, only the HA information is listed (e.g., A/H1pdm09, A/H3).
- The strain name is represented by the location of isolation; in the case of isolation in Japan, the location is written in Japanese kanji, and in the case of isolation in foreign countries, in English.
- To distinguish swine influenza viruses recovered from humans from seasonal influenza viruses, the term “variant virus” is used with a letter “v” added after the subtype name [e.g., A(H3N2)v].

The statistics in this report are based on 1) the data concerning patients and laboratory findings obtained by the National Epidemiological Surveillance of Infectious Diseases undertaken in compliance with the Act on the Prevention of Infectious Diseases and Medical Care for Patients with Infectious Diseases, and 2) other data covering various aspects of infectious diseases. The prefectural and municipal health centers and public health institutes (PHIs), the Department of Environmental Health and Food Safety, the Ministry of Health, Labour and Welfare, and quarantine stations, have provided the above data.

(特集つづき) (THE TOPIC OF THIS MONTH-Continued)

表1. インフルエンザウイルス分離・検出報告数, 2017/18～2020/21シーズン
Table 1. Isolation/detection of influenza viruses during the 2017/18-2020/21 influenza season

型 Type	シーズン* Season*			
	2017/18	2018/19	2019/20	2020/21
A/H1pdm09	1,889 (450)	2,444 (637)	3,802 (1,388)	2 (-)
A/H3	2,319 (1,002)	3,339 (1,396)	95 (38)	4 (-)
A NT	— (15)	4 (10)	2 (4)	— (-)
B/Victoria	113 (31)	476 (130)	508 (242)	— (-)
B/Yamagata	3,486 (938)	37 (9)	3 (1)	— (-)
B NT	6 (52)	6 (6)	— (5)	— (-)
C	40 (17)	1 (2)	14 (4)	1 (-)
合計 Total	7,853 (2,505)	6,307 (2,190)	4,424 (1,682)	7 (-)

A NT: A亜型未同定, B NT: B系統未同定

—報告なし

A NT: A not subtyped, B NT: B lineage not determined


*各シーズン（当年9月～翌年8月）に採取された検体から各地方衛生研究所で分離されたウイルス報告数、()内はウイルスは分離されていないが、遺伝子検出または抗原検出による報告数を別掲

(NESID 病原体検出情報: 2021年9月30日現在報告数)

* Sampling season from September through August in the following year.
(): Nos. in parentheses denote gene or antigen detection without isolation, not included in the total.
[Infectious Agents Surveillance System: as at 30 September 2021 from prefectural and municipal public health institutes (PHIs)]

図2. 都道府県別インフルエンザウイルス分離報告状況, 2020/21シーズン

Figure 2. Isolation of influenza viruses by prefecture in the 2020/21 influenza season



(NESID 病原体検出情報: 2021年9月30日現在報告数)

(Infectious Agents Surveillance System: as at 30 September 2021 from PHIs)

注) IASRのインフルエンザウイルス型、亜型、株名の記載方法は、赤血球凝集素(HA)の分類を調べた情報を主とする場合と、さらにノイラミニダーゼ(NA)の型別まで実施された場合などの違いによるものである。

・N型別まで実施されている場合: A(H1N1)pdm09, A(H3N2), A(H5N1)など

・N型別未実施のものが含まれる場合: A/H1pdm09, A/H3など

・株名については、主に国内の地名は漢字、国外は英語表記(例: B/山形系統, B/Victoria系統など)

・ヒトに感染したブタインフルエンザウイルスはヒトの季節性インフルエンザウイルスと区別するために、variant virusと総称し、亜型の後に“v”を表記: A(H3N2)vなど