

Acute Respiratory Infection Surveillance Weekly Report: Epidemiologic Situational Awareness

Week 24, 2026 (June 8, 2026 – June 14, 2026)

This report aims to systematically review and compile nationwide surveillance data on acute respiratory infections (ARI), and to provide epidemiological information to public health professionals and the general public. Influenza and coronavirus disease 2019 (COVID-19) are reported from ARI sentinel sites consisting of pediatrics and internal medicine departments, while respiratory syncytial virus (RSV) infection, herpangina, pharyngoconjunctival fever, and group A streptococcal pharyngitis are reported from pediatric sentinel sites.

Beginning April 7, 2025 (Week 15), the sentinel selection criteria were revised: Influenza/COVID-19 sentinel sites (approximately 5,000 medical facilities) were replaced by ARI sentinel sites (approximately 3,000 medical facilities), and the number of pediatric sentinel sites was reduced from approximately 3,000 to approximately 2,000. About 10% of those 3,000 sentinel sites send specimens to public health laboratories of each prefecture and are registered as ARI pathogen sentinel sites.

For patient surveillance, data from the most recent week are aggregated as of the compilation date, while data from previous weeks are presented as previously reported, without re-aggregation. For laboratory surveillance, data for all periods are aggregated as of the compilation date. The status of infectious disease activity is interpreted by considering both weekly “trends” and “levels.” Important notes are provided at the end of this report. Please note that the reported numbers are provisional and subject to revision.

Weekly Situation Overview

In week 24 of 2026 (June 8, 2026–June 14, 2026), the number of ARI cases per sentinel site was 47.05 (173,896 cases), representing an increase compared with the previous week. A total of 32 prefectures showed an increase in ARI cases per sentinel site compared with the previous week.

The number of cases reported per sentinel site for each disease was 0.58 for COVID-19, 0.07 for influenza, 2.58 for group A streptococcal pharyngitis, 0.66 for herpangina, 0.60 for pharyngoconjunctival fever, and 0.35 for RSV infection. A total of 12 new hospital admissions due to influenza were reported, representing no change compared with the previous week. 107 new hospital admissions due to COVID-19 were reported, representing an increase of 16 cases from the previous week.

By age group, the highest number of reported cases was observed among individuals aged 10–59 years for influenza and 0–9 years for COVID-19; 1–4 years for RSV infection, pharyngoconjunctival fever, and herpangina and 5–14 years for group A streptococcal pharyngitis.

Among specimens collected in week 24 of 2026 and reported by the time of analysis, 3 specimens tested positive for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), 1 specimen tested positive for RSV, and no specimens tested positive for influenza A virus and influenza B virus.

Contents

Weekly Situation Overview	2
1. Patient Surveillance.....	4
1.1. Nationwide Cases per Sentinel Site	4
1.2. Nationwide Reported Cases by Age Group.....	6
1.3. Cases per Sentinel Site by Prefecture	10
1.4. Nationwide New Hospital Admissions for Influenza and COVID-19.....	23
2. Laboratory Surveillance	24
2.1. Nationwide Reported Cases by Pathogen.....	24
Definition of region.....	29
Interpretive Notes.....	29
References	30
Supplementary information 1. Test results by specimen collection week using fully automated molecular testing systems, such as BioFire FilmArray and BioFire SpotFire.....	31
Supplementary information 2. Weekly cases per sentinel site by prefecture for each disease	32

1. Patient Surveillance

1.1. Nationwide Cases per Sentinel Site


In week 24 of 2026, a total of 3,696 ARI sentinel sites nationwide reported ARI cases. The number of cases per sentinel site was 47.05 (173,896 cases in total) (Figure 1), corresponding to a week-on-week ratio of 1.04 compared with the previous week.

Among reports from ARI sentinel sites, the number of cases per sentinel site was 0.07 for influenza (272 cases) and 0.58 for COVID-19 (2,147 cases) (Figure 1A). The number of reporting sentinel sites was 3,732.

Among reports from pediatric sentinel sites, the number of cases per sentinel site was 0.35 for RSV infection (788 cases), 0.60 for pharyngoconjunctival fever (1,365 cases), 0.66 for herpangina (1,496 cases), and 2.58 for group A streptococcal pharyngitis (5,834 cases) (Figure 1B). The number of reporting pediatric sentinel sites was 2,257.

Regarding recent trends, influenza increased compared with the previous week; COVID-19 increased for 2 consecutive weeks; RSV infection increased compared with the previous week; pharyngoconjunctival fever decreased compared with the previous week; herpangina increased for 5 consecutive weeks; and group A streptococcal pharyngitis remained stable.

Figure 1. Weekly number of ARI cases reported per ARI sentinel site



Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: April 7, 2025 – June 14, 2026).

Figure 1A. Weekly number of influenza and COVID-19 cases reported per ARI sentinel site






Figure 1B. Weekly number of RSV infection, pharyngoconjunctival fever, herpangina, and group A streptococcal pharyngitis cases reported per pediatric sentinel site





Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: April 7, 2025 – June 14, 2026)

Note: The number of cases reported is reproduced in the Infectious Diseases Weekly Report (IDWR) for the corresponding week.


1.2. Nationwide Reported Cases by Age Group

Among ARI cases reported from sentinel sites in week 24 of 2026, the number of reported cases by age group was 122,885 cases among individuals aged 0–9 years (week-on-week ratio: 1.03), 39,250 cases among individuals aged 10–59 years (week-on-week ratio: 1.03), and 11,761 cases among individuals aged 60 years and older (week-on-week ratio: 1.02) (Figure 2).

For trends in reported cases by age group by disease, please refer to Table 1A and Table 1B.

Weekly reported cases by age group for influenza and COVID-19 are shown in Figures 2A and 2B. Among individuals aged 60 years and older, 24 influenza cases and 152 COVID-19 cases were reported among individuals aged 60 years and older; of these, 9 influenza cases and 37 COVID-19 cases were reported among individuals aged 80 years and older.

Figure 2. Weekly reported ARI cases by age group



Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: April 7, 2025 – June 14, 2026)
Note: The number of cases reported is reproduced in the IDWR for the corresponding week.

Figure 2A. Weekly number of reported influenza and COVID-19 cases by age group





Figure 2B. Weekly number of reported cases of RSV infection, pharyngoconjunctival fever, herpangina, and group A streptococcal pharyngitis by age group



Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: April 7, 2025 – June 14, 2026)
 Note: The number of cases reported is reproduced in the IDWR for the corresponding week.

Table 1A. Reported cases and week-on-week ratio (values in parentheses) of influenza and COVID-19 by age group in week 24

Age group	Influenza	COVID-19
0–9 years	97	1,070
	(1.67)	(1.37)

Age group	Influenza	COVID-19
10–59 years	151 (0.94)	925 (1.38)
≥60 years	24 (1.14)	152 (1.11)
Total	272 (1.14)	2,147 (1.35)

Table 1B. Reported cases and week-on-week ratio (values in parentheses) of RSV infection, pharyngoconjunctival fever, herpangina, and group A streptococcal pharyngitis by age group in week 24

Age group	RSV infection	Pharyngoconjunctival fever	Herpangina	Group A streptococcal pharyngitis
0 years	210 (1.11)	139 (0.87)	186 (1.63)	50 (1.14)
1–4 years	546 (1.15)	1,047 (1.02)	1,158 (1.55)	1,869 (0.98)
5–14 years	30 (1.88)	161 (0.84)	147 (1.34)	3,631 (1.02)
≥15 years	2 (0.67)	18 (0.62)	5 (0.83)	284 (0.98)
Total	788 (1.16)	1,365 (0.97)	1,496 (1.53)	5,834 (1.00)

Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: June 8, 2026 – June 14, 2026)

Note: Data for the previous week were referred to the corresponding week's IDWR. Detailed age-specific reported case numbers are available in the IDWR (Category V infectious diseases under sentinel surveillance). When the number of

reported cases in the previous week was zero, the week-on-week ratio is indicated by “-”.

1.3. Cases per Sentinel Site by Prefecture

In week 24 of 2026, the three prefectures with the highest numbers of ARI cases per sentinel site were Gunma, which recorded the highest value at 67.93, followed by Iwate at 66.67, and Saitama at 64.37 (Figure 3A). The number of prefectures in which cases per sentinel site increased compared with the previous week was 32 (Table 2). Across all prefectures, the numbers of cases per sentinel site ranged from 26.05 to 67.93 (Figure 4).

The three prefectures with the highest numbers of cases per sentinel site by disease were Okinawa, Okayama, and Ibaraki for influenza; Miyazaki, Kagoshima, and Ehime for COVID-19; Nagasaki, Saga, and Kagoshima for RSV infection; Shimane, Kagoshima, and Fukuoka for pharyngoconjunctival fever; Oita, Fukuoka, and Kagoshima for herpangina; Ehime, Tottori, and Yamagata for group A streptococcal pharyngitis (Table 3).

Figure 3A. Number of ARI cases reported per ARI sentinel site by prefecture in week 24






Figure 3B. Number of reported cases per sentinel site by prefecture for each infectious disease in week 24


Influenza




COVID-19




RSV infection




Pharyngoconjunctival fever



Herpangina



Group A streptococcal pharyngitis



Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026
(data range: June 8, 2026 – June 14, 2026)

Table 2. Number of ARI cases per sentinel site by prefecture in week 24

Prefecture	Reported number of cases	Number of cases per sentinel	Week-on-week ratio
Hokkaido	6,954	41.89	0.95
Aomori	2,285	44.80	0.97
Iwate	2,800	66.67	1.04
Miyagi	3,447	62.67	1.01
Akita	806	32.24	1.00
Yamagata	2,019	53.13	1.11
Fukushima	2,677	55.77	1.08
Ibaraki	3,718	54.68	1.11
Tochigi	2,780	59.15	0.99
Gunma	3,057	67.93	1.05
Saitama	11,136	64.37	0.98
Chiba	9,631	54.11	1.05
Tokyo	19,155	45.83	1.01
Kanagawa	13,387	56.01	0.98
Niigata	2,061	39.63	0.93
Toyama	2,801	58.35	0.96
Ishikawa	2,202	46.85	0.91
Fukui	1,016	26.05	0.92
Yamanashi	946	27.03	0.96
Nagano	2,757	55.14	1.05
Gifu	1,567	34.82	1.14
Shizuoka	3,107	38.36	0.81
Aichi	8,599	52.75	1.00

Prefecture	Reported number of cases	Number of cases per sentinel	Week-on-week ratio
Mie	2,183	31.64	1.03
Shiga	1,595	40.90	1.09
Kyoto	2,796	44.38	1.07
Osaka	9,564	33.56	1.09
Hyogo	6,921	42.72	1.08
Nara	1,485	35.36	1.06
Wakayama	1,494	33.20	1.06
Tottori	1,224	42.21	1.00
Shimane	846	42.30	1.02
Okayama	2,298	45.96	1.10
Hiroshima	3,890	41.83	1.02
Yamaguchi	3,113	51.03	1.07
Tokushima	908	27.52	1.07
Kagawa	727	31.61	1.16
Ehime	2,219	58.39	1.10
Kochi	1,025	27.70	0.98
Fukuoka	6,155	50.45	1.08
Saga	1,153	48.04	1.18
Nagasaki	2,875	57.50	1.12
Kumamoto	4,041	56.92	1.14
Oita	2,624	45.24	1.11
Miyazaki	993	35.46	1.13
Kagoshima	2,786	48.88	1.08
Okinawa	2,073	47.11	1.05

Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026
(data range: June 8, 2026 – June 14, 2026)

Notes: Data for the previous week were referred to the corresponding week's IDWR. When the number of reported cases in the previous week was zero, the week-on-week ratio is indicated by “-”.


Table 3. Top three prefectures by cases per sentinel site for each infectious disease in week 24







Infectious diseases	Prefectures		
Influenza	Okinawa (0.77)	Okayama (0.26)	Ibaraki (0.24)
COVID-19	Miyazaki (4.11)	Kagoshima (3.46)	Ehime (2.58)
RSV infection	Nagasaki (2.58)	Saga (1.67)	Kagoshima (1.35)
Pharyngoconjunctival fever	Shimane (1.18)	Kagoshima (1.10)	Fukuoka (1.04)
Herpangina	Oita (3.58)	Fukuoka (2.53)	Kagoshima (2.48)
Group A streptococcal pharyngitis	Ehime (6.57)	Tottori (6.26)	Yamagata (5.00)







Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: June 8, 2026 – June 14, 2026)







Note: When cases per sentinel site were identical, prefectures are listed in ascending order of prefecture code.







Figure 4. Weekly reported ARI cases per sentinel site by prefecture


































Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026
(data range: April 7, 2025 – June 14, 2026)


Note: The number of cases reported is reproduced in the IDWR for the corresponding week.

1.4. Nationwide New Hospital Admissions for Influenza and COVID-19

Trends in the number of new hospital admissions reported from designated sentinel medical facilities in week 24 of 2026 are shown in Figure 5, and the number of reported cases by age group is presented in Table 4. A total of 12 new hospital admissions due to influenza were reported, representing no change compared with the previous week. 107 new hospital admissions due to COVID-19 were reported, representing an increase of 16 cases from the previous week.

For the number of cases and trends in each age group, please refer to Table 4.

Figure 5. Weekly number of new hospital admissions due to influenza and COVID-19 reported by designated sentinel hospitals



Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: April 7, 2025 – June 14, 2026)

Table 4. Number of new hospital admissions and week-on-week ratio (values in parentheses) by age group, reported by designated sentinel hospitals in week 24

Age group	Influenza	COVID-19
0–9 years	3 (0.50)	37 (1.06)
10–59 years	0 (0.00)	18 (1.20)
60–69 years	2 (—)	5 (1.00)
70–79 years	4 (4.00)	18 (2.25)
≥80 years	3 (1.00)	29 (1.04)
Total	12 (1.00)	107 (1.18)

Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: June 8, 2026 – June 14, 2026)

2. Laboratory Surveillance

2.1. Nationwide Reported Cases by Pathogen


Among specimens collected at ARI pathogen sentinel sites in week 24 of 2026 and reported by the time of analysis, a total of 70 specimens were reported. Of these, no specimens were positive for influenza A virus or influenza B virus, 3 specimens were positive for SARS-CoV-2, and 1 specimen was positive for RSV (Figure 6).

The pathogen-specific test positivity was 0.0% for influenza A virus and influenza B virus, 1.4% for RSV, and 4.3% for SARS-CoV-2 (Figure 7).

Specimens collected in week 19 (May 4–10) have mostly been registered with test results at the time of aggregation. For the numbers and the most frequently detected pathogen by region, please refer to Table 5.

Test results by specimen collection week using fully automated genetic testing systems at participating medical institutions are presented in Supplementary information 1. For week 24, 3 specimens of rhinovirus/enterovirus were reported.

Figure 6. Weekly number of detected pathogens based on specimen collection week



Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: April 7, 2025 – June 14, 2026).


Note: Data are aggregated by specimen collection week, not by reporting week. The number of test results reflects the data available at the time of aggregation, so they do not necessarily match the figures published in previous reports. When multiple pathogens are detected from a single specimen, all detected pathogens are counted.

“Rhinovirus/Enterovirus” indicates that either rhinovirus or enterovirus was detected.

“Other pathogens” denotes detection of pathogens not listed in the legend.


For weeks and regions with no detections or no reports, it should be noted that this may indicate either that no pathogens were detected or that tests were not performed, depending on the test items.


Figure 7. Weekly pathogen-specific test positivity based on specimen collection week



Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: April 7, 2025 – June 14, 2026).
Note: The test positivity is calculated using the number of specimens tested for the target pathogen as the denominator: $(\text{number positive} / \text{number tested}) \times 100$. Data are aggregated by specimen collection week, not by reporting week. The number of test results reflects the data available at the time of aggregation, so they do not necessarily match the figures published in previous reports.

Figure 8. Weekly reported cases by pathogen at the national and regional levels by specimen collection week





Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: April 7, 2025 – June 14, 2026).

Note: Data are aggregated by specimen collection week, not by reporting week. The number of test results reflects the data available at the time of aggregation and may not necessarily match figures published in previous reports. When multiple pathogens are detected from a single specimen, all detected pathogens are counted. “Rhinovirus/Enterovirus” indicates that either rhinovirus or enterovirus was detected. “Other pathogens” refers to pathogens not listed in the legend. For weeks and regions with no detections or no reports, it should be noted that this may indicate either that no pathogens were detected or that tests were not performed, depending on the test items.

Table 5. Number of specimens and most frequently detected pathogen by region in week 19 (May 4–10)

Region	Number of specimens	Most frequently detected pathogen
--------	---------------------	-----------------------------------

Region	Number of specimens	Most frequently detected pathogen
Hokkaido & Tohoku	48	Rhinovirus/Enterovirus
Kanto	68	Rhinovirus/Enterovirus
Hokuriku	40	Rhinovirus/Enterovirus
Tokai	5	Rhinovirus/Enterovirus
Kinki	42	Rhinovirus/Enterovirus
Chugoku	60	Rhinovirus/Enterovirus
Shikoku	5	Rhinovirus/Enterovirus
Kyushu & Okinawa	34	Rhinovirus/Enterovirus

Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026

Definition of region

Hokkaido & Tohoku: Hokkaido, Aomori, Iwate, Miyagi, Akita, Yamagata, Fukushima

Kanto: Ibaraki, Tochigi, Gunma, Saitama, Chiba, Tokyo, Kanagawa, Yamanashi, Nagano

Hokuriku: Niigata, Toyama, Ishikawa, Fukui

Tokai: Gifu, Shizuoka, Aichi, Mie

Kinki: Shiga, Kyoto, Osaka, Hyogo, Nara, Wakayama

Chugoku: Tottori, Shimane, Okayama, Hiroshima, Yamaguchi

Shikoku: Tokushima, Kagawa, Ehime, Kochi

Kyushu & Okinawa: Fukuoka, Saga, Nagasaki, Kumamoto, Oita, Miyazaki, Kagoshima, Okinawa

Interpretive Notes

Sentinel definitions and the composition of reporting sites changed on 7 April 2025 (week 15). Time-series comparisons across this date must be interpreted with caution. Figures in the original report demarcate this change.

Reporting tends to decrease during certain holiday periods, such as the year-end/New Year holidays (around weeks 52–1), Golden Week (around week 18), the Obon holidays (around week 33), and Silver Week (around week 39). The specific weeks may vary by year depending on the arrangement of public holidays and weekends.

“Cases per sentinel site” reflect both disease activity and care-seeking/reporting behavior; shifts in the denominator (participation, holidays) can influence observed levels.

Counts are provisional and subject to backfill due to delayed reporting and data correction.

Laboratory surveillance data shown for all weeks reflect the information available at the time of compilation. Testing items for specimens collected may vary, depending on municipalities or regional public health laboratories. In addition, because the time required for testing and reporting differs among these laboratories, the number of pathogen detections for a given specimen collection week may be delayed or later revised. Thus, aggregated values should be considered provisional.

References

- Infectious Diseases Weekly Report (IDWR)
<https://id-info.jihs.go.jp/en/surveillance/idwr/index.html>
- Infectious Agents Surveillance Report (IASR)
<https://id-info.jihs.go.jp/en/surveillance/iasr/index.html>
- Japan Institute for Health Security (JIHS) The Infectious Disease Information Website
<https://id-info.jihs.go.jp/en/>
- Ministry of Health, Labour and Welfare website [Japanese]
 - Acute Respiratory Infection (ARI)
<https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/kenkou/kekkaku-kansenshou19/ari.html>
 - Influenza
https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/kenkou_iryuu/kenkou/kekkaku-kansenshou/infuenza/index.html
 - COVID-19
https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000164708_00001.html
 - RSV infection
https://www.mhlw.go.jp/bunya/kenkou/kekkaku-kansenshou19/rs_qa.html
 - Pharyngoconjunctival fever
https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/kenkou_iryuu/kenkou/kekkaku-kansenshou/pcf.html
 - Herpangina
https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/kenkou_iryuu/kenkou/kekkaku-kansenshou/herpangina.html
- Graphical Overview of Infectious Diseases
https://www.jihs.go.jp/content10/030/en_Dashboard.html
- Genomic surveillance of SARS-CoV-2 (including quarantine specimens and specimens from incoming travelers)[Japanese]
<https://id-info.jihs.go.jp/surveillance/iasr/45/532/article/030/index.html>
- Variants of SARS-CoV-2 [Japanese]
<https://id-info.jihs.go.jp/relevant-information/covid-19/variants/index.html>

Supplementary information 1. Test results by specimen collection week using fully automated molecular testing systems, such as BioFire FilmArray and BioFire SpotFire

Test results from pathogen testing conducted at medical institutions equipped with fully automated genetic testing systems are presented below. These data are collected through voluntary participation of selected medical institutions and are used for monitoring purposes.

Pathogen	Week 19	Week 20	Week 21	Week 22	Week 23	Week 24
Influenza A/H1	0	0	0	0	0	0
Influenza A/H1pdm09	0	0	0	0	0	0
Influenza A/H3	0	0	0	0	0	0
Influenza A virus (subtype unknown)	0	0	0	0	0	0
Influenza B virus	0	0	0	0	0	0
SARS-CoV-2	1	1	1	0	0	0
RSV	1	3	2	0	0	0
Parainfluenza virus 1	1	1	0	0	0	0
Parainfluenza virus 2	0	0	0	0	0	0
Parainfluenza virus 3	0	0	0	1	0	0
Parainfluenza virus 4	0	0	0	0	0	0
Parainfluenza virus (type unknown)	0	0	0	0	0	0
Rhinovirus/Enterovirus	4	7	9	5	8	3
Human metapneumovirus	2	4	6	5	2	0
Adenovirus	1	0	2	0	3	0
Coronavirus HKU1	0	0	0	0	0	0
Coronavirus NL63	0	0	0	0	0	0
Coronavirus 229E	2	0	0	0	0	0
Coronavirus OC43	1	0	0	0	0	0
Bordetella pertussis	0	0	0	0	0	0
Bordetella parapertussis	0	0	0	0	0	0

Pathogen	Week 19	Week 20	Week 21	Week 22	Week 23	Week 24
Chlamydia pneumoniae	0	0	0	0	0	0
Mycoplasma pneumoniae	0	0	0	0	1	0

Source: Infectious Disease Surveillance in Japan; data as of June 17, 2026 (data range: May 4, 2026 to June 14, 2026)


Note: As reporting is based on voluntary participation by medical institutions, the number of reported cases should be interpreted as reference values. A total of 15 medical institutions participated between weeks 19–24.

Rhinovirus/Enterovirus indicates detection of either rhinovirus or enterovirus.


Records labeled only as “cov” or “flu” are excluded from this table.

Supplementary information 2. Weekly cases per sentinel site by prefecture for each disease


Hokkaido




Aomori




Iwate




Miyagi




Akita




Yamagata




Fukushima




Ibaraki




Tochigi




Gunma




Saitama




Chiba




Tokyo




Kanagawa




Niigata




Toyama




Ishikawa




Fukui




Yamanashi




Nagano




Gifu




Shizuoka




Aichi




Mie




Shiga




Kyoto




Osaka




Hyogo




Nara




Wakayama




Tottori




Shimane




Okayama




Hiroshima




Yamaguchi




Tokushima




Kagawa




Ehime




Kochi




Fukuoka




Saga




Nagasaki




Kumamoto




Oita




Miyazaki



Kagoshima



Okinawa



Data source: Infectious Disease Surveillance in Japan; data as of June 17, 2026
(data range: April 7, 2025 – June 14, 2026)